



Rec'd PCT/PTO 20 DEC 2004
PCT/CH 03 / 00401

REC'D 08 JUL 2003

WIPO

PCT



INVESTOR IN PEOPLE

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.


In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated 2 May 2003



23 JUN 2007 10:14:49
P01/770000.00-0214344.4

THE PATENT OFFICE

21 JUN 2007

NEWPORT

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

Request for grant of a patent

See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference

30037 GB

2. Patent application number

(The Patent Office will fill in this part)

0214344.4

21 JUN 2007

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Givaudan SA
Chemin de la Parfumerie 5
1214 Vernier
Switzerland

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

8408031001

4. Title of the invention

New odorant compounds

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

CIT Givaudan UK Ltd.
Magna House
76-80 Church Street
Staines, Middlesex TW18 4XR
England

Patents ADP number (if you know it)

8408056001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

Yes

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

- Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form

Description 16

Claim(s) 6

Abstract 1

Drawing(s) 1

0. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) 1

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

12. Name and daytime telephone number of person to contact in the United Kingdom

Colin Brown
Office time

Tel. No.:

01 784417721

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
b) Write your answers in capital letters using black ink or you may type them.
c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

d) If you have answered 'Yes' Patent Form 7/77 will need to be filed.

New odorant compounds

This invention relates to novel compounds having woody, vetiver and patchouli-like odour notes. This invention relates furthermore to a method for their production and to
5 flavour and fragrance compositions containing them.

Compounds having woody, vetiver and patchouli-like odour notes are described in the literature, for example the class of sesquiterpenes that naturally occur in essential oils and which can be isolated by water-steam distillation of a plant or parts of a plant. This
10 process is very cost intensive and the quality and the odour as well as the flavour characteristics of the isolated compounds may vary with the climate and the origin of the plant. Thus, there is an ongoing demand in the fragrance and flavour industry for new compounds imparting, enhancing, or improving woody, vetiver and patchouli-like notes.


15

Tricyclic sesquiterpenes, in particular patchoulol and derivatives thereof have been disclosed in the U.S. Patent No. 4,011,269. The compounds described therein develop odoriferous notes the character of which is reminiscent of that of patchouli oil.

20

We have now found a novel class of compounds having much sought after woody, vetiver and patchouli-like odour notes and which may be produced from synthetic starting materials.

In a first aspect the invention provides a compound of the formula I



25

wherein

R^1 , R^4 , R^6 and R^7 are independently hydrogen, methyl or ethyl;

R^2 and R^3 are independently hydrogen, or C_{1-5} alkyl, e.g. methyl, ethyl, or linear or branched propyl, butyl, or pentyl; or

R^2 and R^3 together with the carbon atom to which they are attached form a 5- or 6-membered cycloalkyl ring;

5 R^5 is hydrogen, or C_{1-4} alkyl, e.g. methyl, ethyl or linear or branched propyl;

R^6 is hydrogen, or branched lower C_{3-7} alkyl, e.g. isopropyl, tert. butyl;

R^9 is hydrogen, methyl, ethyl, or branched lower C_{3-7} alkyl, e.g. isopropyl, tert. butyl;

R^{10} is ethyl or propyl;

R^{11} is C_{1-4} alkyl, e.g. methyl, ethyl, or linear or branched propyl or butyl;

10 R^{12} is hydroxy;

R^{13} is hydrogen, or C_{1-4} alkyl, e.g. methyl ethyl, or linear or branched propyl or butyl; or

R^{12} and R^{13} together with the carbon atom to which they are attached form a carbonyl group;

the dashed line represents either a C-C single bond or no bond; and

15

a) when C5 and C8 are connected by single bond and C9 and C6 are connected by a single bond, C9 and C5 are not connected by a bond,

$n=1$,

R^7 , R^8 are hydrogen, and

20 R^9 is hydrogen, methyl or ethyl; or

b) when C5 and C8 are connected by a single bond and C9 and C6 are connected by a single bond, C9 and C5 are not connected,

$n=0$,

25 R^7 , R^8 is hydrogen,

R^9 is a branched lower C_{3-7} alkyl; or

c) when C5 and C8 are not connected by a bond, C9 and C5 are connected by a single bond,

30 R^7 is hydrogen, methyl or ethyl,

R^8 is a branched lower C_{3-7} alkyl, or


R^7 and R^8 together with the carbon atoms to which they are attached form a 5- or 6-membered cycloalkyl ring,

$n = 0$, and

35 the bond between C6 and C8 may be a single bond or a double bond.

1703-11-11

Preferred compounds are those of the formulae Ia, Ib and Ic



5

wherein

R^1 , R^4 , R^6 , and R^{16} are independently hydrogen, methyl or ethyl;

R^7 and R^{14} are independently hydrogen, methyl or ethyl; or,

10 R^7 and R^{14} together with the carbon atoms to which they are attached form a 5- or 6-membered cycloalkyl ring;

R^2 and R^3 are independently hydrogen, or C_{1-5} alkyl, e.g. methyl, ethyl, or linear or branched propyl, butyl, or pentyl; or,

15 R^2 and R^3 together with the carbon atom to which they are attached form a 5- or 6-membered cycloalkyl ring;

R^5 is hydrogen, or C_{1-4} alkyl, e.g. methyl, ethyl, or linear or branched propyl;

R^{15} is C_{1-4} alkyl, e.g. methyl, ethyl, or linear or branched propyl;

R^{12} is hydroxy;

R^{13} is hydrogen or C_{1-4} alkyl, e.g. methyl, ethyl, or linear or branched propyl; or

20 R^{12} and R^{13} together with the carbon atom to which they are attached form a carbonyl group; and

in formula Ic the bond between C6 and C8 may be a single bond, or the dotted line together with the bond between C6 and C8 may represent a double bond .

The compounds according to the present invention contain one or more chiral centres and as such may exist as a mixture of stereoisomers, or they may be resolved as isomerically pure forms. Resolving stereoisomers adds to the complexity of manufacture and purification of these compounds and so it is preferred to use the compounds as mixtures of their stereoisomers simply for economic reasons. However, if it is desired to prepare individual stereoisomers, this may be achieved according to methodology known in the art, e.g. preparative HPLC and GC or by stereoselective syntheses.

Particular preferred compounds of formula Ia are 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one, 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one, 1,3,3,5,7,8,8-Heptamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one, 3,3,5,7,8,8-Hexamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one, 3,3,5,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one, 5,7,8,8-Tetramethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one, and 5,6,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-ol.

A particular preferred compound of formula Ib is 1-Isopropyl-3,3,5-trimethyl-tricyclo[3.2.1.0^{2,7}]octan-6-one.

Particular preferred compounds of formula Ic are 5-Isopropyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one, 5-Isopropyl-1,3-dimethyl-bicyclo[3.2.1]octan-2-one, 5-tert-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one, 5-sec-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-ene-2-one, 5-Isopropyl-3-methyl-bicyclo[3.2.1]oct-3-ene-2-one, 5,7-Diisopropyl-3-methyl-bicyclo[3.2.1]oct-3-en-2-one, 5-Isopropyl-3,7,7-trimethyl-bicyclo[3.2.1]oct-3-en-2-one, and 1,3,5-Trimethyl-1,5,6,7,8,8a-hexahydro-1,4a-ethano-naphthalen-2-one.

In another aspect the invention provides flavour and fragrance compositions comprising a compound of formula I or mixtures thereof, more particularly compounds of formula Ia, Ib, or Ic or mixtures thereof. Particular preferred are compositions that comprise at least one compound of formula Ia and one compound of formula Ic.

In addition, the compounds may be used in combination with other known odourant molecules selected from the extensive range of natural and synthetic molecules currently available, such as essential oil, alcohols, aldehydes and ketones, ether and

acetals, ester and lactones, macrocycles and heterocycles, and/or in admixture with one or more ingredients or excipients conventionally used in conjunction with odourants in fragrance compositions, for example carrier materials, and other auxiliary agents commonly used in the art.

5

In one embodiment, the compounds of the present invention may be used in fragrance applications, e.g. in any field of fine and functionary perfumery, such as perfumes, household products, laundry products, body care products and cosmetics.

10 In another embodiment, the compounds of the present invention may be used in flavour applications and are particularly useful in modifying for example strawberry and raspberry flavours but also brown flavours. They may be used in herbal mixtures and teas. The compounds of the present invention are also well suited for example in mouthwash applications.

15

In flavourant applications, the compounds of the present invention may be present in compositions in amounts ranging from 0.001 to 5 % by weight of a flavour composition, more preferably from 0.01 to 0.5 % by weight. The compounds according to the present invention may be used for herbal flavour compositions, strawberry and raspberry
20 compositions, brown flavour compositions, or tea compositions.

25

When used in fragrance applications, compounds of the present invention can be employed in wide ranging amounts depending upon the specific application and on the nature and quantity of other odourant ingredients, that may be for example, from about
0.001 to about 20 weight percent. In one embodiment compounds may be employed in a fabric softener comprising in amount of about 0.001 to 0.05 weight percent. In another
embodiment compounds of the present invention may be in alcoholic solution in amounts of about 0.1 to 20 weight percent, more preferably between about 0.1 and 5 weight percent.


30

However, these values should not be limiting on the present invention, since the experienced perfumer and flavourist may also achieve effects or may create novel accords with lower or higher concentrations.

The compounds according to the present invention may be prepared according to a process wherein appropriately substituted cyclohexenones are reacted with allyl bromide or allyl chloride under reaction conditions well known to the person skilled in the art (Bull. Chem. Soc. Jpn., 2298 – 2303 (1993)). The resulting alkylated cyclohexenones (formula II as shown below) may be converted in the presence of ethyl aluminium-dichloride or methyl aluminium-dichloride to provide compounds of formula I wherein R^{12} and R^{13} taken together represents an oxygen atom, as illustrated by compounds of formula Ia, Ib, and Ic in scheme 1. The conditions under which such Lewis acid catalyzed reactions may proceed is described for example by Snider et al. in the Journal of Am. Chem. Soc. 1980, 102, 5872 – 5880 which is herein incorporated by reference.

The resulting carbonyl group at C1 may be reduced and/or alkylated to give further compounds of formula I. Similarly, if there is a double bond at C6 and C8 this can be reduced in a known manner to give still further compounds of formula I.

Scheme1:



Compounds of formula II may also be prepared by alkylation of appropriately substituted phenols by reaction of the phenol with a metal hydride and an alkenylchloride (Greuter, H. et al. (1977) Helv. Chim. Acta, 60, 1701), followed by hydrogenation.

Whether a compound of formula Ia, Ib, Ic, or a mixture thereof is formed depends on the substituent pattern R^2 , R^3 , and R^{16} of the alkylated cyclohexenone (II). A compound of formula Ic is formed as a main product if R^2 , R^3 , and R^{16} of formula II are hydrogen. A mixture of compounds of formula Ia and Ib as main product is performed if R^2 and R^3 of formula II at the same time are not hydrogen. A mixture of compounds of formula Ia and Ic as main product is formed if R^2 and R^3 of formula II are hydrogen and R^{16} of formula II is not hydrogen. The compounds are useful in flavour and/or fragrance compositions as mixtures, however, should one wish to use the compounds in pure form, they can be separated easily by purification processes, such as HPLC or preparative GC, according to the methodology known in the art.

The term "main product", as used herein with reference to single compounds, refers to a product comprising at least 50% by weight of that compound, more preferably more than 75% by weight, most preferably more than 90% by weight. When this term is used in relation to a mixture of compounds, e.g. one compound of formula Ia and one compound of formula Ic, it refers to a product comprising at least 50% by weight of this mixture, more particular more than 75% by weight, most particular more than 90% by weight.

The conversion of compounds of formula II for a selective preparation of compounds of formula Ia of the present invention may also be performed by photochemical induction. Surprisingly we found that compounds of formula Ia may be formed by photochemically induced intramolecular [2+2] cycloaddition of compounds of formula II. For the photochemical induction a Hg-lamp may be used for a time period of about 1 to 15 hours. However, the induction time may depend on the solvent used and on additives such as sensitisers and Lewis acids. Preferred solvents are methanol, ethanol and iso propanol.

Thus, in another aspect the invention refers to a method of preparing compounds of formula Ia by photochemically induced cycloaddition.

Further particulars as to reaction conditions are provided in the examples.

There now follows a series of examples that illustrate the invention.

Example 1:a) 2-Methyl-6-(3-methyl-but-2-enyl)-cyclohex-2-enone

To a solution of LDA (prepared from BuLi, 1.6 M in hexane, 75 ml, 0.12mol and diisopropylamine, 12.2g, 0.12mol) in THF (50ml) was added 2-methyl-cyclohex-2-enone
 5 (11.0g, 0.1mol) at -78°C . The mixture was stirred for 1h at -78°C and prenyl bromide 17.88g, 0.12mol) was added. The mixture was stirred over night, while the temperature was allowed to rise to room temperature. MTBE (50 ml) and sat. NH_4Cl were added, the organic phase was separated and washed with brine, dried (MgSO_4) and concentrated in vacuo. The residue was distilled in a Kugelrohr oven to yield 9.8g (55%) of a slightly
 10 yellow oil.

$^1\text{H-NMR}$ (400 MHz, CDCl_3): 6.82 (bs, 1H), 5.14-5.09 (m, 1H), 2.56-2.48 (m, 1H), 2.37-2.25 (m, 3H), 2.13-2.03 (m, 2H), 1.77 (s, 3H), 1.70 (s, 3H), 1.75-1.68 (m, 1H), 1.61 (s, 3H) ppm. $^{13}\text{C-NMR}$ (100MHz, CDCl_3): 201.6 (s), 144.3 (d), 135.1 (s), 133.0 (s), 121.8 (d), 46.9 (d), 27.8 (t), 27.7 (t), 25.7 (q), 25.0 (t), 17.6 (q), 16.0 (q) ppm. GC/MS (EI): 178
 15 (M^+ , 30), 168 (25), 123 (37), 110 (100), 95 (63), 83 (33), 69 (26), 53 (34), 41 (71), 39 (44). IR (ATR): 2966s, 2924s, 1672vs, 1451s, 1377s, 1181m, 1088m, 836m cm^{-1} .

b) 2,6-dimethyl-6-(3-methyl-but-2-enyl)-cyclohex-2-enone

Sodium hydride (60%, 85 g, 2.13 mol) was added portionwise to a solution of 2,6-
 20 dimethylphenol (250g, 2.05 mol) in 2L of toluene at $10-15^{\circ}\text{C}$. The resulting suspension was stirred for 45 min. The mixture was cooled to 5°C , and prenyl chloride (262g, 2.13 mol, 85%) was added during 1.5 h keeping the temperature at 5°C . The mixture was then stirred for further 2h at $10-15^{\circ}\text{C}$. Methanol (1L) and palladium (2.5g, 10% on charcoal) was added and the gray suspension was hydrogenated at 0.3 bar
 25 overpressure, keeping the temperature at $20-22^{\circ}\text{C}$ (ice bath). The suspension was then filtered through a pad of celite. The yellow filtrate was washed with water (0.5L), aqueous sodium hydroxide (0.5L) and brine (0.5L), dried (MgSO_4) and concentrated *in vacuo*. The residue was distilled over a 5cm Vigreux column to yield 318g (81%, bp $78-82^{\circ}\text{C}/0.05\text{Torr}$) of a colorless oil.

30 Odor description: fruity, grapefruit, minty, bergamot

$^1\text{H-NMR}$ (400MHz, CDCl_3): 6.62 (bs, 1H, 3-H), 5.06-5.11 (m, 1H, 2'-H), 2.34-2.28 (m, 2H, 4-H), 2.25-2.14 (m, 2H, 1'-H), 1.91 (dt, $J_{5a,5b} = 13.6 \text{ Hz}$, $J_{5a,4} = 6.1 \text{ Hz}$, 1H, 5a-H), 1.76 (s, 3H, 2- CH_3), 1.77-1.70 (m, 1H, 5b-H), 1.70 (s, 3H, 4'-H), 1.59 (s, 3H, 3'- CH_3),

1.05 (s, 3H, 6-CH₃) ppm. GC/MS (EI): 192 (M⁺, 16), 124 (100), 109 (74), 82 (31), 69 (40), 41 (57). IR (ATR): 2965s, 2922s, 1667vs, 1449m, 1376m, 1033m cm⁻¹.

Example 2: 5-Isopropyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one

5 To a solution of 2,6-dimethyl-6-(3-methyl-but-2-enyl)-cyclohex-2-enone (5.00 g, 26.04 mmol) in toluene (40 ml) was added dropwise neat EtAlCl₂ (97%, 1.5 eq., 4.96 g, 39.06 mmol). During the addition, the temperature was kept below 10°C. The brown mixture was kept at room temperature over night and was then poured on icecold saturated NH₄Cl. The mixture was extracted with MTBE, washed with brine, dried (MgSO₄) and
10 concentrated in vacuo. The residue was distilled bulb to bulb to yield 4.50 g (90%) of an colorless oil.

Odor description: woody, patchouli, vetiver, hesperidic

¹H-NMR (400MHz, CDCl₃): 6.86 (bs, 1H, 4-H), 1.81 (dt, *J* = 11.0, 2.2 Hz, 1H, 8-H_a), 1.78-1.72 (m, 2H, 6-H_a, 7-H_a), 1.75 (d, *J* = 1.6 Hz, 3-CH₃), 1.69 (sept, *J* = 6.8 Hz, 1H, 5-CH(CH₃)₂), 1.64-1.53 (m, 2H, 6-H_b, 7-H_b), 1.37 (dd, *J* = 11.0, 2.2 Hz, 1H, 8-H_b), 1.24 (s, 3H, 1-CH₃), 1.00 (d, *J* = 6.8Hz, 3H, 5-CH(CH₃)CH₃), 0.93 (d, *J* = 6.8 Hz, 3H, CH(CH₃)CH₃) ppm. ¹³C-NMR (100MHz, CDCl₃) : 205.2 (s, C-2), 151.7 (d, C-4), 133.1 (s, C-3), 52.4 (s, C-1), 51.8 (s, C-5), 50.7 (t, C-8), 34.8 (t, C-6), 34.8, (d, 5-CH(CH₃)₂), 33.3 (t, C-7), 20.7 (q, 1-CH₃), 19.0, 18.4 (2q, CH-(CH₃)₂), 15.6 (q, 3-CH₃) ppm. GC/MS
20 (EI): 192 (M⁺, 24), 177 (18), 149 (52), 136 (33), 121 (95), 110 (100), 91(46), 77 (39), 41 (41). IR (ATR) : 2959s, 2867m, 1674vs, 1446m, 1362m, 1331m, 1030s cm⁻¹.

Example 3: 5-Isopropyl-1,3-dimethyl-bicyclo[3.2.1]octan-2-one

Prepared by hydrogenation of 5-isopropyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one.

25 Mixture of 2 isomers (ratio of 5/1), main isomer: ¹H-NMR (400MHz, CDCl₃): 2.46-2.39 (m, 1H), 2.13-2.06 (m, 1H), 1.88-1.43 (m, 8H), 1.29-1.26 (m, 1H), 1.16 (s, 3H, 1-CH₃), 1.11 (d, *J* = 7.2 Hz, 3H, 3-CH₃), 1.04-0.97 (m, 1H), 0.92 (d, *J* = 6.8 Hz, 5-CH(CH₃)CH₃), 0.88 (d, *J* = 6.8 Hz, 5-CH(CH₃)CH₃) ppm. ¹³C-NMR (100MHz, CDCl₃): 219.6 (s, C-2), 52.7 (s), 46.3 (s) 44.7 (t), 38.7 (d), 37.9 (t), 37.5 (d), 36.3 (t), 36.4 (t), 20.4 (q), 18.4 (q),
30 17.4 (q), 16.5 (q) ppm. GC/MS (EI): 194 (M⁺, 14), 151 (100), 133 (11), 123 (48), 93 (29), 81 (83), 69 (19), 41 (30). IR (ATR): 2958s, 2868m, 1709vs, 1458s, 1369m, 999m cm⁻¹.

Odor description : woody, ambery, ionone

Example 4 - 12:

The following compounds were prepared according to the synthetic procedure of Example 2 from the correspondingly substituted materials and purified by chromatography where indicated.

5

Mixture of 5-tert-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one and 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one

Prepared as a mixture and purified by chromatography.

a) 5-tert-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one

10 Odor description: woody, patchouli, vetiver

¹H-NMR (400MHz, CDCl₃): 7.03 (bs, 1H, 4-H), 1.99 (ddd, *J* = 16 Hz, 12 Hz, 5.6 Hz, 1H, 8-H_a), 1.75 (d, *J* = 1.6 Hz, 3-H, 3-CH₃), 1.75-1.53 (m, 5H), 1.47-1.40 (m, 1H), 1.25 (s, 3H, 1-CH₃), 0.98 (s, 9H, 5-C(CH₃)₃) ppm. ¹³C-NMR (100MHz, CDCl₃): 204 (s, C-2), 152.6 (d, C-4), 132.3 (s, C-3), 54.2 (s), 51.9 (s), 46.9 (t, C-8), 33.5 (t), 32.9 (s, C(CH₃)₃), 30.7 (t), 25.9 (q, C(CH₃)₃), 20.7 (q, 1-CH₃), 15.5 (q, 3-CH₃) ppm. GC/MS (EI): 206 (M⁺, 6), 191 (8), 149 (10), 135 (24), 124 (99), 110 (100) 91 (32), 77 (17), 57 (28), 41 (35). IR (ATR): 2961s, 2868m, 1673vs, 1467m, 1446m, 1365m, 1238m, 1030m, 879w cm⁻¹.

15

b) 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one

20 Odor description: woody, patchouli

¹H-NMR (400MHz, CDCl₃): 2.19 (bs, 1H, 2-H), 1.78 (dd, *J*_{9a,9b} = 12.8 Hz, *J* = 2.0 Hz, 1H, 9-H_a), 1.76-1.50 (m, 4H, 3,4-H), 1.55 (d, *J*_{9a,9b} = 12.8 Hz, 1H, 9-H_b), 1.05 (s, 3H, 8-(CH₃)_a), 0.98 (s, 3H, 5-CH₃), 0.97 (s, 3H, 7-CH₃), 0.96 (s, 3H, 1-CH₃), 0.63 (s, 3H, 8-(CH₃)_b) ppm. ¹³C-NMR (100MHz, CDCl₃): 221.1 (s, C-6), 57.0 (s, C-7), 46.6 (d, C-2), 45.5 (s, C-8), 43.7 (s, C-5), 42.6 (t, C-9), 41.8 (s, C-1), 38.6 (t, C-4), 21.0 (q, 8-(CH₃)_a), 19.7 (2q, C-5, 8-(CH₃)_b), 18.1 (q, 1-CH₃), 10.8 (q, 7-CH₃) ppm. GC/MS (EI): 206 (M⁺, 6), 191 (10), 124 (100), 109 (27), 97 (76), 69 (22) 55 (17), 41 (31). IR (ATR): 2926m, 2862m, 1711s, 1446m, 1373m, 1004m, 761w cm⁻¹.

25

30 5-sec-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one

Odor description: woody, vetiver, cedar, olibanum

Mixture of 2 isomers : ¹H-NMR (400MHz, CDCl₃): 6.86, 6.83 (2bs, 1H, 4-H), 1.95-1.51 (m, 7H), 1.75 (s, 3H, 3-CH₃), 1.40-1.32 (m, 2H), 1.23 (s, 3H, 1-CH₃), 1.00-0.90 (m, 6H) ppm. GC/MS (EI): 206 (M⁺, 8), 177 (17), 149 (41), 124 (100), 121 (82), 110 (98), 91

(42), 77 (32), 41 (43). IR (ATR): 2961s, 2865m, 1674vs, 1446s, 1364s, 1222m, 1034s cm^{-1} .

5-Isopropyl-3-methyl-bicyclo[3.2.1]oct-3-ene-2-one

5 Odor description: ambery, ciste, woody

$^1\text{H-NMR}$ (400MHz, CDCl_3): 6.88 (bs, 1H, 4-H), 2.96 (dd, $J_{1,7a} = 7.4 \text{ Hz}$, $J_{1,8b} = 4.6 \text{ Hz}$, 1H, 1-H), 2.15 (m, 1H, 7- H_a), 1.92 (bd, $J_{8a,8b} = 11.2 \text{ Hz}$, 1H, 8- H_a), 1.74 (d, $J = 1.2 \text{ Hz}$, 3H, 3- CH_3), 1.72 (sept, $J = 6.8 \text{ Hz}$, 1H, 5- $\text{CH}(\text{CH}_3)_2$), 1.72-1.58 (m, 2H, 6- H_a , 7- H_b), 1.52-1.45 (m, 1H, 6- H_b), 1.42 (ddd, $J_{8b,8a} = 11.2 \text{ Hz}$, $J_{8b,1} = 4.6 \text{ Hz}$, $J = 2.0 \text{ Hz}$, 8- H_b),
 10 1.03 (d, $J = 6.8 \text{ Hz}$, 3H, 5- $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.95 (d, $J = 6.8 \text{ Hz}$, 3H, 5- $\text{CH}(\text{CH}_3)\text{CH}_3$) ppm.
 $^{13}\text{C-NMR}$ (100MHz, CDCl_3): 204.1 (s, C-2), 152.1 (d, C-4), 133.2 (s, C-3), 51.2 (s, C-5), 50.2 (d, C-1), 43.3 (t, C-8), 34.6 (d, 5- $\text{CH}(\text{CH}_3)_2$), 33.8 (t, C-6), 25.5 (t, C-7), 19.0, 18.5 (2q, 5- $\text{CH}(\text{CH}_3)_2$), 15.2 (q, 3- CH_3) ppm. GC/MS (EI): 178 (M^+ , 49), 163 (61), 135 (50), 123 (21), 107 (100), 91 (59), 79 (46), 77 (39), 67 (23), 41 (47). IR (ATR): 2957s, 2871m,
 15 1677vs, 1447m, 1358s, 1053m, 1018m, 918s cm^{-1} .

5,7-Diisopropyl-3-methyl-bicyclo[3.2.1]oct-3-en-2-one

Odor description: woody, elemi

Mixture of the endo/exo-isomers in a ratio of 2/1; main (endo) isomer: $^1\text{H-NMR}$
 20 (400MHz, CDCl_3): 6.97 (bs, 1H, 4-H), 3.06 (dd, $J = 6.2, 4.4 \text{ Hz}$, 1H, 1-H), 2.03-1.96 (m, 2H), 1.87 (dd, $J = 12.8, 10.4 \text{ Hz}$, 1H), 1.76-1.51 (m, 3H), 1.72 (d, $J = 1.6 \text{ Hz}$, 3H, 3- CH_3), 1.39-1.33 (m, 1H), 1.00 (d, $J = 6.8 \text{ Hz}$, 3H), 0.96 (d, $J = 6.0 \text{ Hz}$, 3H), 0.91 (d, $J = 6.8 \text{ Hz}$, 3H), 0.85 (d, $J = 6.0 \text{ Hz}$, 3H) ppm. $^{13}\text{C-NMR}$ (100MHz, CDCl_3): 203.0 (s, C-2), 153.9 (d, C-4), 134.0 (s, C-3), 54.4 (d, C-1), 50.2 (s, C-5), 48.5 (d, C-7), 44.4 (t, C-8),
 25 39.1 (t, C-6), 34.7 (d), 32.4 (d), 22.2 (q), 21.7 (q), 18.6 (q), 18.3 (q), 15.0 (q, 3- CH_3) ppm. GC/MS (EI): 220 (M^+ , 17), 205 (8), 177 (22), 151 (28), 135 (32), 121 (30), 109 (100), 107 (50), 91 (47), 77 (29), 69 (35), 41 (52). IR (ATR): 2956 s, 2872m, 1673vs, 1467m, 1365s, 973m, 903m cm^{-1} .

30 5-Isopropyl-3,7,7-trimethyl-bicyclo[3.2.1]oct-3-en-2-one

Prepared according to Example 2 with 2,5,5-Trimethyl-6-(3-methyl-but-2-enyl)-cyclohex-2-enone as starting material.

$^1\text{H-NMR}$ (400MHz, CDCl_3): 6.88 (bs, 1H, 4-H), 2.52 (d, $J_{1,8b} = 4.4 \text{ Hz}$, 1H, 1-H), 1.98 (bd, $J_{8a,8b} = 11.2 \text{ Hz}$, 1H, 8- H_a), 1.80 (ddd, $J_{8b,8a} = 11.2 \text{ Hz}$, $J_{8b,1} = 4.4 \text{ Hz}$, $J = 2.0 \text{ Hz}$,

1H, 8-H_b), 1.73 (s, 3H, 3-CH₃), 1.66 (sept., $J = 7.0$ Hz, 1H, 5-CH(CH₃)₂), 1.53 (s, 2H, 6-H), 1.19 (s, 3H, 7-(CH₃)_a), 0.97 (d, $J = 7.0$ Hz, 3H, 5-CH(CH₃)CH₃), 0.91 (d, $J = 7.0$ Hz, 3H, 5-CH(CH₃)CH₃), 0.90 (s, 3H, 7-(CH₃)_b) ppm. ¹³C-NMR (100MHz, CDCl₃): 203.7 (s, C-2), 152.2 (d, C-4), 133.3 (s, C-3), 62.4 (d, C-1), 51.7 (s, C-5), 48.5 (t, C-8), 42.6 (t, C-6), 38.8 (s, C-7), 35.0 (d, 5-CH(CH₃)₂), 32.2 (q, 7-(CH₃)_a), 27.5 (q, 7-(CH₃)_b), 18.5 (q), 18.2 (q) ppm. GC/MS (EI): 206 (M⁺, 35), 191 (26), 163 (25), 135 (100), 121 (50), 107 (98), 91 (73), 77 (53), 69 (29), 55 (34), 41 (81). IR (ATR): 2957s, 2870m, 1676vs, 1466m, 1359m, 1048w, 885w cm⁻¹.

Odor description: woody, resin

10

1,3,5-Trimethyl-1,5,6,7,8,8a-hexahydro-1,4a-ethano-naphthalen-2-one

Odor description: woody, patchouli, vetiver, cedar

Mixture of two isomers; ¹H-NMR (400MHz, CDCl₃, (1S^{*},4aR^{*},5S^{*},8aR^{*})-isomer,

derived from HMQC): 6.63 (bs, 1H, 4-H), 1.77 (d, $J = 1.6$ Hz, 3H, 3-CH₃), 1.78-1.68 (m,

15 3H, 8-H_a, 9-H_a, 10-H_a), 1.63-1.53 (m, 2H, 9-H_b, 10-H_b), 1.54-1.47 (m, 1H, 6-H_a), 1.50-1.44 (m, 1H, 7-H_a), 1.45-1.35 (m, 1H, 5-H), 1.42-1.36 (m, 1H, 8a-H), 1.22-1.14 (m, 1H, 8-H_b), 1.16 (s, 3H, 1-CH₃), 1.08-0.94 (m, 2H, 6-H_b, 7-H_b), 1.03 (d, $J = 6.8$ Hz, 3H, 5-CH₃) ppm. ¹³C-NMR (100MHz, CDCl₃): 205.0 (s, C-2), 147.2 (d, C-4), 134.7 (s, C-3), 61.3 (d, C-8a), 54.5 (s, C-1), 50.7 (s, C-4a), 41.2 (d, C-5), 34.8 (t, C-9), 33.2 (t, C-10), 20 31.8 (t, C-6), 25.9 (t, C-8), 21.8 (t, C-7), 18.8 (q, 1-CH₃), 16.6 (q, 5-CH₃), 15.6 (q, 3-CH₃) ppm. MS(mixture) (EI): 218 (M⁺, 96), 203 (12), 190 (7), 175 (13), 161 (7), 147 (17), 124 (100), 105 (20), 95 (58), 82 (22), 67 (8), 55 (13), 41 (24).

1,3,3,5,7,8,8-Heptamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one

25 Prepared as a mixture and purified by chromatography.

Odor description: patchouli, vetiver, woody

¹H-NMR (400MHz, CDCl₃): 1.96 (bs, 1H, 2-H), 1.79 (dd, $J_{9a,9b} = 13.2$ Hz, $J_{9a,4b} = 2.8$ Hz,

1H, 9-H_a), 1.66 (d, $J_{9b,9a} = 13.2$ Hz, 1H, 9-H_b), 1.59 (dd, $J_{4a,4b} = 13.2$ Hz, $J = 1.0$ Hz, 1H,

4-H_a), 1.48 (dd, $J_{4b,4a} = 13.2$ Hz, $J_{4b,9a} = 2.8$ Hz, 1H, 4-H_b), 1.08 (s, 3H, 5-CH₃), 1.04 (s,

30 3H, 8-(CH₃)_a), 1.04 (s, 3H, 7-CH₃), 1.03 (s, 3H, 3-(CH₃)_a), 0.97 (s, 3H, 1-CH₃), 0.83 (s, 3H, 3-(CH₃)_b), 0.58 (s, 3H, 8-(CH₃)_b) ppm. ¹³C-NMR (100MHz, CDCl₃): 221.3 (s, C-6), 58.7 (d, C-2), 56.1 (s), 53.6 (t, C-4), 44.4 (s), 44.3 (s), 41.6 (s), 44.1 (t, C-9), 31.3 (s), 31.3 (2q, 3-(CH₃)_{a,b}), 21.2 (q, 8-(CH₃)_a), 20.3 (q, 5-CH₃), 19.8 (q, 1-CH₃), 18.9 (q, 8-(CH₃)_b), 13.4 (q, 7-CH₃) ppm. GC/MS (EI): 234 (M⁺, 28), 219 (20), 163 (15), 152 (43),

137 (62), 121 (49), 97 (199), 83 (18), 69 (25), 57 (72), 41 (69). IR (ATR): 2959m, 2919m, 2865m, 1712s, 1452m, 1374m, 1005m, 951m, 886w cm^{-1} .

3,3,5,7,8,8-Hexamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one

5 Prepared as a mixture and purified by chromatography.

Odor description: ambery, woody, pine resin, Grisalva

¹H-NMR (400MHz, CDCl_3): 2.32 (t, $J = 6.0$ Hz, 1H, 1-H), 2.21 (d, $J = 6.0$ Hz, 1H, 2-H), 1.90 (dd, $J = 13.2$ Hz, 6.0 Hz, 1H, 9- H_a), 1.74 (dd, $J = 13.2$ Hz, 2.6 Hz, 1H), 1.57 (d, $J = 13.2$ Hz, 1-H), 1.48 (dd, $J = 13.2$, 2.6 Hz, 1H), 1.17 (s, 3H), 1.04 (s, 3H), 1.00 (s, 3H), 0.97 (s, 3H), 0.84 (s, 3H), 0.64 (s, 3H) ppm. ¹³C-NMR (100MHz, CDCl_3): 220.9 (s), 57.3 (s), 53.3 (t), 52.6 (d), 44.0 (s), 42.3 (s), 39.6 (d), 33.4 (t), 30.9 (s), 30.5 (q), 30.3 (q), 23.9 (q), 20.1 (q), 20.0 (q), 13.4 (q) ppm. GC/MS (EI): 220 (M^+ , 6), 205 (5), 164 (64), 152 (37), 137 (199), 123 (19), 91 (14), 69 (26), 55 (17), 41 (46). IR (ATR): 2956m, 2924m, 2866w, 1711s, 1451m, 1373m, 1012m, 956m, cm^{-1} .

15

Mixture of 3,3,5,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one and 1-Isopropyl-3,3,5-trimethyl-tricyclo[3.2.1.0^{2,7}]octan-6-one

Prepared as a mixture and purified by chromatography.

a) 3,3,5,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one

20 Odor description: woody, patchouli, cedar, camphoraceous

¹H-NMR (400MHz, CDCl_3): 2.61 (t, $J = 5.0$ Hz, 1H, 7-H), 2.46 (bt, $J = 5.6$ Hz, 1H, 2-H), 2.34 (bq, $J = 5.6$ Hz, 1H, 1-H), 1.92 (dd, $J_{9a,9b} = 13.2$ Hz, $J_{9a,1} = 5.6$ Hz, 1H, 9- H_a), 1.73 ($J_{9b,9a} = 13.2$ Hz, $J = 2.8$ Hz, 1H, 9- H_b), 1.58 (d, $J_{4a,4b} = 13.6$ Hz, 1H, 4- H_a), 1.49 (dd, $J_{4b,4a} = 13.6$ Hz, $J = 2.8$ Hz, 1H, 4- H_b), 1.32 (s, 3H, 8- $(\text{CH}_3)_a$), 1.02 (s, 3H, 3- $(\text{CH}_3)_a$), 0.97 (s, 3H, 5- CH_3), 0.82 (s, 3H, 3- $(\text{CH}_3)_b$), 0.75 (s, 3H, 8- $(\text{CH}_3)_b$) ppm. ¹³C-NMR (100MHz, d_6 -acetone): 219.0 (s, C-6), 57.6 (d, C-7), 53.2 (t, C-4), 47.2 (d, C-2), 44.3 (s, C-3), 40.7 (d, C-1), 40.6 (s, C-5), 33.6 (t, C-9), 29.9 (q, 3- $(\text{CH}_3)_a$), 29.8 (q, 3- $(\text{CH}_3)_b$), 29.8 (s, C-8), 27.1 (q, 8- $(\text{CH}_3)_a$), 21.6 (q, 8- $(\text{CH}_3)_b$), 19.8 (q, 5- CH_3) ppm. GC/MS (EI): 206 (M^+ , 17), 191, (15), 150 (29), 138 (145), 123 (64), 107 (48), 83 (100), 55 (29), 41 (48). IR (ATR): 2953s, 2866m, 1716vs, 1459m, 1373m, 1105m, 1002w, 931w, 898w cm^{-1} .

30

b) 1-Isopropyl-3,3,5-trimethyl-tricyclo[3.2.1.0^{2,7}]octan-6-one

Odor description: woody, cedar, veriver, fruity, patchouli

¹H-NMR (400MHz, C₆D₆): 1.62 (d, $J = 8.3$ Hz, 1H, 7-H), 1.59 (dd, $J_{8a,8b} = 11.2$ Hz, $J = 2.4$ Hz, 1H, 8-H_a), 1.52 (d, $J_{8b,8a} = 11.2$ Hz, 8-H_b), 1.45 (dd, $J_{4a,4b} = 13.6$ Hz, $J = 2.2$ Hz, 1H, 4-H_a), 1.26 (d, $J_{4b,4a} = 13.6$ Hz, 1H, 4-H_b), 1.21-1.11 (m, 2H, 2-H, 1-CH(CH₃)₂), 1.01 (s, 3H, 5-CH₃), 0.99 (s, 3H, 3-(CH₃)_a), 0.96 (s, 3H, 3-(CH₃)_b), 0.83 (d, $J = 6.8$ Hz, 3H, CH(CH₃)_a(CH₃)_b), 0.80 (d, $J = 6.8$ Hz, 3H, CH(CH₃)_a(CH₃)_b) ppm. ¹³C-NMR (100MHz, C₆D₆): 212.6 (s, C-6), 52.2 (t, C-4), 46.0 (d, C-2), 42.7 (s, C-5), 40.2 (s, C-1), 34.3 (t, C-8), 34.1 (d, C-7), 32.0 (2q, 3-(CH₃)_{a,b}), 31.9 (d, 1-CH(CH₃)₂), 29.2 (s, C-3), 18.9 (q, 5-CH₃), 18.8 (q, 1-CH(CH₃)_a(CH₃)_b), 18.7 (q, 1-CH(CH₃)_a(CH₃)_b) ppm. GC/MS (EI): 206 (M⁺, 41), 191 (37), 151 (15), 135 (36), 109 (100), 91 (38), 77 (24), 55 (21), 41 (43). IR (ATR): 2957m, 2926m, 2867m, 1725s, 1462m, 1317m, 1171m, 917m, 866 m, 834m cm⁻¹.

Example 13: 5,7,8,8-Tetramethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one

A solution of 2,6-dimethyl-6-(3-methyl-but-2-enyl)-cyclohex-2-enone (10.0g, 52.1 mmol) in methanol (250 ml) was irradiated using a Hg-lamp during 3h. The solvent was evaporated in vacuo and the residue distilled in a Kugelrohr oven to yield 5.0 g of a colorless oil.

Odor description: patchouli, woody, camporaceous

¹H-NMR (400MHz, CDCl₃): 2.54 (m, 1H, 2-H), 2.23 (t, $J = 5.8$ Hz, 1H, 1-H), 1.88 (dd, $J_{9a,9b} = 12.8$ Hz, $J = 5.6$ Hz, 1H, 9-H_a), 1.78 (d, $J_{9b,9a} = 12.8$ Hz, 1H, 9-H_b), 1.88-1.52 (m, 4H, 3,4-H), 1.17 (s, 3H, 8-(CH₃)_a), 1.00 (s, 3H, 5-CH₃), 0.99 (s, 3H, 7-CH₃), 0.66 (s, 3H, 8-(CH₃)_b) ppm. ¹³C-NMR (100MHz, C₆D₆): 217.6 (s, C-6), 58.1 (s, C-7), 42.7 (2s, C-5,8), 41.3 (d, C-1), 40.7 (d, C-2), 38.1 (t, C-4), 34.4 (t, C-9), 23.5 (q, 8-(CH₃)_a), 20.4 (q, 8-(CH₃)_b), 20.2 (q, 5-CH₃), 17.5 (t, C-3), 11.1 (s, 7-CH₃) ppm. GC/MS (EI): 192 (M⁺, 14), 177 (12), 149 (8), 124 (100), 109 (48), 93 (15), 82 (20), 69 (34), 53 (18), 41 (52). IR (ATR): 2923m, 2863m, 1710s, 1448m, 1375m, 1068m, 1020m, 1000m, 790w cm⁻¹.

Example 14: 5,6,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-ol

Prepared from 5,7,8,8-tetramethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one by reaction with methyl magnesium chloride.

Odor description: patchouli, woody, camphoraceous

¹H-NMR (400MHz, C₆D₆): 2.23 (dd, $J_{9a,9b} = 13.2$ Hz, $J_{9a,4a} = 3.2$ Hz, 1H, 9-H_a), 2.12-2.09 (m, 1H, 2-H), 1.95 (dd, $J_{1,9b} = 6.8$ Hz, $J_{1,2} = 6.0$ Hz, 1H, 1-H), 1.62-1.49 (m, 3H, 4-H_a, 3-H_{a,b}), 1.43 (dd, $J_{9b,3a} = 13.2$ Hz, $J_{9b,1} = 6.8$ Hz, 1H, 9-H_b), 1.37 (s, 3H, 8-(CH₃)_a), 1.26-1.18 (m, 1H, 4-H_b), 1.21 (s, 3H, 8-(CH₃)_b), 1.04 (s, 3H, 5-CH₃), 0.97 (s, 3H, 6-CH₃), 0.93 (s, 3H, 7-CH₃) ppm.

3H, 7-CH₃) ppm. ¹³C-NMR (100MHz, C₆D₆): 79.6 (s, C-6), 50.4 (s, C-7), 42.1 (s, C-8), 41.7 (d, C-1), 38.7 (d, C-2), 36.9 (s, C-5), 36.5 (t, C-9), 35.8 (t, C-4), 27.9 (q, 8-(CH₃)_a), 24.9 (q, 6-CH₃), 21.6 (q, 8-(CH₃)_b), 20.9 (q, 5-CH₃), 18.0 (t, C-3), 12.8 (q, 7-CH₃) ppm.
 GC/MS (EI): 208 (M⁺, 1), 190 (22), 175 (28), 162 (9), 147 (48), 121 (73), 98 (59), 83
 5 (37), 69 (19), 55 (31), 43 (100), 41 (43). IR (ATR): 3502br., 2947vs, 2902vs, 2869s, 1457s, 1371s, 1207m, 1110s, 1072s, 1045s, 922vs cm⁻¹.

Example 15:

A perfume for a shower gel with a woody-floral character

10

parts per weight

	Cedryl acetate	5
	Citronellyl acetate	2
	Linalyl acetate	20
15	Agrumex	20
	Pheyl ethyl alcohol	40
	Amyl cinnamyl aldehyde	140
	Ambrettolide	5
	Ambrofix	4
20	Bois Cedre ess. Virginie	20
	Bois Gaiac ess.	10
	Damascenone (10% in DPG)	6
	Dipropylene glycol	10
	Eucalyptus ess.	24
25	Galaxolide 50 BB	381
	Hedione	80
	Javanol	2
	Lilial	30
	Linalool synt.	20
30	Mandarine ess.	30
	Moxalone	40
	N 112	1
	Okoumal	5
	Orange terpenes dist.	60
35	Rose abs. Turquie	2


	Rose artess abs.	10
	Rose oxide (10% in DPG)	5
	Super muguet	10
	Vanilline (10% in DPG)	8
5	1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0 ^{2,7}]nonan-6-one	10

		1000

10 In this composition 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one enhances and harmonizes the woody part. It underlines the patchouli aspect but gives also a new modern woody character. The compound blends well with the floral part of the perfume and provides volume without giving a heavy impression.

Claims

1. A compound of formula I



wherein

R^1 , R^4 , R^6 and R^7 are independently hydrogen, methyl or ethyl;

R^2 and R^3 are independently hydrogen, or C_{1-5} alkyl; or

R^2 and R^3 together with the carbon atom to which they are attached form a 5- or 6-membered cycloalkyl ring;

R^5 is hydrogen, or C_{1-4} alkyl;

R^8 is hydrogen, or branched lower C_{3-7} alkyl;

R^9 is hydrogen, methyl, ethyl, or branched lower C_{3-7} alkyl;

R^{10} is ethyl or propyl;

R^{11} is C_{1-4} alkyl;

R^{12} is hydroxy;

R^{13} is hydrogen, or C_{1-4} alkyl; or

R^{12} and R^{13} together with the carbon atom to which they are attached form a carbonyl group;

the dashed line represents either a C-C single bond or no bond; and

a) when C5 and C8 are connected by a single bond and C9 and C6 are connected by a single bond, C9 and C5 are not connected by a bond,


$n=1$,

R^7 , R^8 are hydrogen, and

R^9 is hydrogen, methyl or ethyl; or

- b) when C5 and C8 are connected by a single bond and C9 and C6 are connected by a single bond, C9 and C5 are not connected,
 $n=0$,
 R^7, R^8 is hydrogen,
 R^9 is a branched lower C_{3-7} alkyl; or
- c) when C5 and C8 are not connected by a bond, C9 and C5 are connected by a single bond,
 R^7 is hydrogen, methyl or ethyl,
 R^8 is a branched lower C_{3-7} alkyl, or
 R^7 and R^8 together with the carbon atoms to which they are attached form a 5- or 6-membered cycloalkyl ring,
 $n = 0$, and
the bond between C6 and C8 may be a single bond or a double bond.


2. A compound according to claim 1 having a formula Ia



wherein

- R^1, R^4, R^6, R^{14} and R^{16} are independently hydrogen, methyl or ethyl;
 R^2 and R^3 are independently hydrogen, or C_{1-5} alkyl; or,
 R^2 and R^3 together with the carbon atom to which they are attached form a 5- or 6-membered cycloalkyl ring;
 R^5 is hydrogen, or C_{1-4} alkyl;
 R^{15} is C_{1-4} alkyl;
 R^{12} is hydroxy;
 R^{13} is hydrogen or C_{1-4} alkyl; or
 R^{12} and R^{13} together with the carbon atom to which they are attached form a carbonyl group.

3. A compound according to claim 1 of formula Ib,



wherein

R^1 , R^4 , R^6 , R^{14} and R^{16} are independently hydrogen, methyl or ethyl;

R^2 and R^3 are independently hydrogen, or C_{1-5} alkyl; or,

R^2 and R^3 together with the carbon atom to which they are attached form a 5- or 6-membered cycloalkyl ring;

R^5 is hydrogen, or C_{1-4} alkyl;


R^{15} is C_{1-4} alkyl;

R^{12} is hydroxy;

R^{13} is hydrogen or C_{1-4} alkyl; or

R^{12} and R^{13} together with the carbon atom to which they are attached form a carbonyl group.

4. A compound according to claim 1 of formula Ic,



wherein

R^1 , R^4 , R^6 , R^{14} and R^{16} are independently hydrogen, methyl or ethyl;

R^5 is hydrogen, or C_{1-4} alkyl;

R^7 and R^{14} are independently hydrogen, methyl or ethyl; or,

R^7 and R^{14} together with the carbon atoms to which they are attached form a 5- or 6-membered cycloalkyl ring;

R^{15} is C_{1-4} alkyl;

R^{12} is hydroxy;

R^{13} is hydrogen or C_{1-4} alkyl; or


R^{12} and R^{13} together with the carbon atom to which they are attached form a carbonyl group; and

the bond between C6 and C8 may be a single bond;


or the dotted line together with the bond between C6 and C8 may represent a double bond.

5. A compound according to claim 1 selected from the group consisting of 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one; 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one; 1,3,3,5,7,8,8-Heptamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one; 3,3,5,7,8,8-Hexamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one; 3,3,5,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one; 5,7,8,8-Tetramethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one; 1-Isopropyl-3,3,5-trimethyl-tricyclo[3.2.1.0^{2,7}]octan-6-one; 5-Isopropyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one; 5-Isopropyl-1,3-dimethyl-bicyclo[3.2.1]octan-2-one; 5-tert-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one; 5-sec-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-ene-2-one; 5-Isopropyl-3-methyl-bicyclo[3.2.1]oct-3-ene-2-one; 5,7-Diisopropyl-3-methyl-bicyclo[3.2.1]oct-3-en-2-one; 5-Isopropyl-3,7,7-trimethyl-bicyclo[3.2.1]oct-3-en-2-one; 1,3,5-Trimethyl-1,5,6,7,8,8a-hexahydro-1,4a-ethano-naphthalen-2-one; and 5,6,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-ol.
6. A flavour or fragrance composition comprising a compound as defined according to one of the preceding claims.
7. A flavour or fragrance composition according to claim 6 comprising at least one compound is selected from the group of compounds of formula Ia as defined in claim 2 and at least one compound is selected from the group of compounds of formula Ic as defined in claim 4.
8. A flavour or fragrance composition according to claim 7 comprising 5-tert-Butyl-1,3-dimethyl-bicyclo[3.2.1]oct-3-en-2-one and 1,5,7,8,8-Pentamethyl-tricyclo[3.3.1.0^{2,7}]nonan-6-one.

9. The use of a compound as defined in one of the claims 1 to 5 in fragrance and flavour applications.
10. The use of a compound according to claim 9 selected in perfumes, household products, laundry products, body care products, and cosmetics.
11. The use according to claim 9 and claim 10 wherein a compound is provided in an amount from 0.001 to 20% by weight.
12. A process of preparing a compound of the formula I as defined in claim 1



comprising the step of reacting a compound of formula II with ethyl aluminium dichloride or methyl aluminium dichloride



wherein

R^1 , R^4 , and R^6 are independently hydrogen, methyl or ethyl;

R^2 and R^3 are independently hydrogen, or C_{1-5} alkyl; or

R^2 and R^3 together with the carbon atom to which they are attached form a 5- or 6-membered cycloalkyl ring;

R^5 is hydrogen, or C_{1-4} alkyl;


R^7 and R^{14} are independently hydrogen, methyl or ethyl; or

R^7 and R^{14} together with the carbon atoms to which they are attached form a 5- or 6-membered cycloalkane ring;


R^{16} is hydrogen, or lower branched C_{3-7} alkyl,

and optionally followed by the step of reduction and/or alkylation of the carbonyl group at C1.

13. A process of preparing a compound of the general formula Ic



comprising the step of converting a compound of formula II by photochemical induction



wherein

R^2 , R^3 , and R^{16} are hydrogen;

R^1 , R^4 and R^6 are independently hydrogen, methyl or ethyl;

R^7 and R^{14} are independently hydrogen, methyl or ethyl; or

R^7 and R^{14} together with the carbon atoms to which they are attached form a 5- or 6-membered cycloalkane ring;

R^5 is hydrogen, linear or branched C_{1-4} alkyl;


R^{15} is linear or branched C_{1-4} alkyl; and

and optionally followed by the step of hydrogenation across the double bond at C6 and C8, and

optionally followed by the step of reduction and/or alkylation of the carbonyl group at C1.

Abstract

Novel compounds of formula I and their use in flavour and fragrance compositions.



R¹ to R¹³ have the meaning as described in the specification.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.